

# Linear programming model for solution of matrix game with payoffs trapezoidal intuitionistic fuzzy number

*Darunee Hunwisai*<sup>1</sup>; *Poom Kumam*<sup>2</sup>

CompAMa Vol.5, No.1, pp.7-30, 2017 - Accepted September 2, 2016

## Abstract

In this work, we considered two-person zero-sum games with fuzzy payoffs and matrix games with payoffs of trapezoidal intuitionistic fuzzy numbers (TrIFNs). The concepts of TrIFNs and their arithmetic operations were used. The cut-set based method for matrix games with payoffs of TrIFNs was also considered. Compute the interval-type value of any alfa-constrategies by simplex method for linear programming. The proposed method is illustrated with a numerical example.

**Keywords:** Intuitionistic fuzzy set, matrix game, linear programming.

## 1 Introduction

The concept of an intuitionistic fuzzy set was proposed by Atanassov in 1986 [1]. This concept referred to the reflect of the relation among "1 minus the degree of membership", "the degree of non-membership" and "the degree of hesitation". The intuitionistic fuzzy set was rasterized by the degree of

---

<sup>1</sup>Department of Mathematics, King Mongkut's University of Technology Thonburi, 126 Pracha-Uthit Road, Bang Mod, Thrung Khru, Bangkok 10140 Thailand, (darunee.3790@mail.kmutt.ac.th).

<sup>2</sup>Theoretical and Computational Science Center (TaCS), Science Laboratory Building, Faculty of Science, King Mongkut's University of Technology Thonburi (KMUTT), 126 Pracha-Uthit Road, Bang Mod, Thrung Khru, Bangkok 10140 Thailand, (poom.kumam@mail.kmutt.ac.th).

membership and the degree of non-membership. The intuitionistic fuzzy set had more abundant and flexible than the fuzzy set with uncertain information. Ishibuchi and Tanaka [2], Chanas and Kuchta [3], studies multiobjective programming in optimization of interval objective functions, solving interval-valued objective optimization problems. More specifically, interval-valued objective optimization problems could be converted into bi-objective mathematical programming models. Then, the bi-objective mathematical programming models were also solved using the existing methods of multi-objective programming. Cevikel and Ahlatçoglu [4] found fuzzy payoffs and fuzzy goals of two-person zero-sum games. The payoff matrix with elements was represented as a fuzzy number. For any pair of the strategies, a player received a payoff that meant as a fuzzy number. Nan *et al* [5], focused a lexicographic method for matrix games with payoffs of triangular intuitionistic fuzzy numbers. Moreover Nan *et al* [6] also introduced a new ranking method based on the value and used to solving matrix games with payoffs on trapezoidal intuitionistic fuzzy numbers (TrIFNs) fuzzy goals. Hence, there were also found the use of linear programming method for solving matrix games [7]-[8]. Aggarwal *et al* [9], solved the matrix games with I-fuzzy payoffs: Pareto-optimal security strategies approach. Therefore, in this work, two-person zero-sum games with fuzzy payoffs, matrix games with payoffs of TrIFNs were considered. The concepts of TrIFNs and their arithmetic operations were used. The matrix games of the cut-set based method with payoffs of TrIFNs was aimed. The auxiliary linear programming models were computed the interval-type value of any  $\alpha$ -constrategies. This paper is organized as follows. In section 2, the definition, operations of TrIFNs and a methodology for matrix games with payoffs TrIFN were focused. In section 3, matrix games with payoffs of TrIFNs were formulated. In section 4, an application to voting share problem was reported. In the last section, section 5, conclusion was summerized.

## 2 Mathematical Preliminaries

In this section, we summarize some basic concept intuitionistic fuzzy set by Atanassov [1], notation, definition and operation of trapezoidal intuitionistic fuzzy number which are used throughout the paper.

## 2.1 Some definitions of TrIFNs

**Definition 1.** [1] Let  $X$  be a nonempty set of the universe. If there are two mapping on the set  $X$ :

$$\begin{aligned}\mu_{\tilde{A}} &: X \rightarrow [0, 1] \\ x &\mapsto \mu_{\tilde{A}}(x)\end{aligned}$$

and

$$\begin{aligned}\nu_{\tilde{A}} &: X \rightarrow [0, 1] \\ x &\mapsto \nu_{\tilde{A}}(x)\end{aligned}$$

with the condition  $0 \leq \mu_{\tilde{A}}(x) + \nu_{\tilde{A}}(x) \leq 1$ . The  $\mu_{\tilde{A}}$  and  $\nu_{\tilde{A}}$  are called determining and intuitionistic fuzzy set  $\tilde{A}$  on the universal set  $X$ , denote by  $\{\langle x; \mu_{\tilde{A}}(x), \nu_{\tilde{A}}(x) \rangle | x \in X\}$  we called  $\mu_{\tilde{A}}$  and  $\nu_{\tilde{A}}$  are membership function and nonmembership function of  $\tilde{A}$ , respectively.  $\mu_{\tilde{A}}(x)$  and  $\nu_{\tilde{A}}(x)$  are called the membership degree and nonmembership degree of an element  $x$  belonging to  $\tilde{A} \subseteq X$ , respectively.  $F(X)$  is called the set of the intuitionistic fuzzy set on the universal set  $X$ .

**Definition 2.** A TrIFN  $\tilde{A} = \langle (l, c, d, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  is a special intuitionistic fuzzy set on the real number set  $\mathbb{R}$ , whose membership and nonmembership functions are defined as follows:

$$\mu_{\tilde{A}}(x) = \begin{cases} 0 & \text{if } x < l \\ t_{\tilde{A}}(x - l)/(c - l) & \text{if } l \leq x < c \\ t_{\tilde{A}} & \text{if } c \leq x \leq d \\ t_{\tilde{A}}(r - x)/(r - d) & \text{if } d < x \leq r \\ 0 & \text{if } x > r \end{cases} \quad (1)$$

and

$$\nu_{\tilde{A}}(x) = \begin{cases} 1 & \text{if } x < l \\ [c - x + z_{\tilde{A}}(x - l)]/(c - l) & \text{if } l \leq x < c \\ z_{\tilde{A}} & \text{if } c \leq x \leq d \\ [x - d + z_{\tilde{A}}(r - x)]/(r - d) & \text{if } d < x \leq r \\ 1 & \text{if } x > r \end{cases} \quad (2)$$

respectively, where  $l \leq c \leq d \leq r$ , the values  $t_{\tilde{A}}$  and  $z_{\tilde{A}}$  are maximum membership degree and minimum nonmembership degree of  $\tilde{A}$ , respectively,




Fig. 1: A TrIFN  $\tilde{A} = \langle (l, c, d, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$

such that they satisfy the following condition:  $t_{\tilde{A}} \in [0, 1]$ ,  $z_{\tilde{A}} \in [0, 1]$  and  $t_{\tilde{A}} + z_{\tilde{A}} \in [0, 1]$ .

Let

$$\pi_{\tilde{A}}(x) = 1 - \mu_{\tilde{A}}(x) - \nu_{\tilde{A}}(x) \quad (3)$$

$\pi_{\tilde{A}}(x)$  is called the hesitancy degree of an element  $x \in \tilde{A}$ . It is the degree of indeterminacy membership of the element  $x$  to  $\tilde{A}$ .

From Definition 2, it is obvious that  $\mu_{\tilde{A}}(x) + \nu_{\tilde{A}}(x) = 1$  for any  $x \in \mathbb{R}$  if  $t_{\tilde{A}} = 1$  and  $z_{\tilde{A}} = 0$ . Hence, the TrIFN  $\tilde{A} = \langle (l, c, d, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  degenerates to  $\tilde{A} = \langle (l, c, d, r); 1, 0 \rangle$ , which is a trapezoidal fuzzy number [10]. Therefore, the concept of the TrIFN is generalization of that of the trapezoidal fuzzy number.

From  $\tilde{A} = \langle (l, c, d, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  if  $c = d = p$  then  $\tilde{A} = \langle (l, p, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  that is  $\tilde{A} = \langle (l, p, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  is a triangular intuitionistic fuzzy number (TIFN), which is particular case of TrIFN. Likewise to algebraic operations of TIFN and TrIFN are defined as follows.

**Definition 3.** Let  $\tilde{A} = \langle (l_1, c_1, d_1, r_1); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  and  $\tilde{B} = \langle (l_2, c_2, d_2, r_2); t_{\tilde{B}}, z_{\tilde{B}} \rangle$  be two TrIFNs with  $t_{\tilde{A}} \neq t_{\tilde{B}}$  and  $z_{\tilde{A}} \neq z_{\tilde{B}}$ ,  $\gamma \neq 0$  be any real number. Then, the algebraic operations of TrIFNs are defined as follows:

$$\tilde{A} + \tilde{B} = \langle (l_1 + l_2, c_1 + c_2, d_1 + d_2, r_1 + r_2); t_{\tilde{A}} \wedge t_{\tilde{B}}, z_{\tilde{A}} \vee z_{\tilde{B}} \rangle, \quad (4)$$

$$\tilde{A} - \tilde{B} = \langle (l_1 - r_2, c_1 - d_2, d_1 - c_2, r_1 - l_2); t_{\tilde{A}} \wedge t_{\tilde{B}}, z_{\tilde{A}} \vee z_{\tilde{B}} \rangle, \quad (5)$$

$$\tilde{A}\tilde{B} = \begin{cases} \langle (l_1 l_2, c_1 c_2, d_1 d_2, r_1 r_2); t_{\tilde{A}} \wedge t_{\tilde{B}}, z_{\tilde{A}} \vee z_{\tilde{B}} \rangle & \text{if } \tilde{A} > 0, \tilde{B} > 0 \\ \langle (l_1 r_2, c_1 d_2, d_1 c_2, r_1 l_2); t_{\tilde{A}} \wedge t_{\tilde{B}}, z_{\tilde{A}} \vee z_{\tilde{B}} \rangle & \text{if } \tilde{A} < 0, \tilde{B} > 0 \\ \langle (r_1 r_2, d_1 d_2, c_1 c_2, l_1 l_2); t_{\tilde{A}} \wedge t_{\tilde{B}}, z_{\tilde{A}} \vee z_{\tilde{B}} \rangle & \text{if } \tilde{A} < 0, \tilde{B} < 0, \end{cases} \quad (6)$$

$$\tilde{A}/\tilde{B} = \begin{cases} \langle (l_1/r_2, c_1/d_2, d_1/c_2, r_1/l_2); t_{\tilde{A}} \wedge t_{\tilde{B}}, z_{\tilde{A}} \vee z_{\tilde{B}} \rangle & \text{if } \tilde{A} > 0, \tilde{B} > 0 \\ \langle (r_1/r_2, d_1/d_2, c_1/c_2, l_1/l_2); t_{\tilde{A}} \wedge t_{\tilde{B}}, z_{\tilde{A}} \vee z_{\tilde{B}} \rangle & \text{if } \tilde{A} < 0, \tilde{B} > 0 \\ \langle (r_1/l_2, d_1/c_2, c_1/d_2, l_1/r_2); t_{\tilde{A}} \wedge t_{\tilde{B}}, z_{\tilde{A}} \vee z_{\tilde{B}} \rangle & \text{if } \tilde{A} < 0, \tilde{B} < 0, \end{cases} \quad (7)$$

$$\gamma/\tilde{A} = \begin{cases} \langle (\gamma l_1, \gamma c_1, \gamma d_1, \gamma r_1); t_{\tilde{A}}, z_{\tilde{A}} \rangle & \text{if } \gamma > 0 \\ \langle (\gamma r_1, \gamma d_1, \gamma c_1, \gamma l_1); t_{\tilde{A}}, z_{\tilde{A}} \rangle & \text{if } \gamma < 0 \end{cases} \quad (8)$$

and

$$\tilde{A}^{-1} = \langle (1/r_1, 1/d_1, 1/c_1, 1/l_1); t_{\tilde{A}}, z_{\tilde{A}} \rangle \quad \text{if } \tilde{A} \neq 0 \quad (9)$$

where the symbols  $\wedge$  is the minimum operator and  $\vee$  is the maximum operator.

## 2.2 A methodology for matrix games with payoffs TrIFN

Let  $S_1 = \{\rho_1, \rho_2, \dots, \rho_m\}$  and  $S_2 = \{\tau_1, \tau_2, \dots, \tau_n\}$  be sets of pure strategies for players I and II, respectively. The vector  $x = (x_1, x_2, \dots, x_m)^T$  is mixed strategies for player I where  $x_i$  ( $i = 1, 2, \dots, m$ ) is probability in player I. The set of mixed strategies for player I is represented by  $X = \{x \mid \sum_{i=1}^m x_i = 1, x_i \geq 0 \ (i = 1, 2, \dots, m)\}$ .

Similarly, the vector  $y = (y_1, y_2, \dots, y_n)^T$  is mixed strategies for player II where  $y_j$  ( $j = 1, 2, \dots, n$ ) is probability in player II. The set of mixed strategies for player II is  $Y = \{y \mid \sum_{j=1}^n y_j = 1, y_j \geq 0 \ (j = 1, 2, \dots, n)\}$ . Assume that

the payoff of players I is expressed with an TrIFN

$$\tilde{D}_{ij} = \langle (l_{ij}, c_{ij}, d_{ij}, r_{ij}); t_{\tilde{D}_{ij}}, z_{\tilde{D}_{ij}} \rangle,$$

where  $l_{ij} \leq c_{ij} \leq d_{ij} \leq r_{ij}$ ,  $t_{\tilde{D}_{ij}} \in [0, 1]$  and  $z_{\tilde{D}_{ij}} \in [0, 1]$  ( $i = 1, 2, \dots, m; j = 1, 2, \dots, n$ ). Therefore, the payoff player I at all  $m \times n$  pure strategy situations can be concisely expressed in the matrix format as follows:

$$\tilde{D} = \begin{bmatrix} \tilde{D}_{11} & \tilde{D}_{12} & \cdots & \tilde{D}_{1n} \\ \tilde{D}_{21} & \tilde{D}_{22} & \cdots & \tilde{D}_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ \tilde{D}_{m1} & \tilde{D}_{m2} & \cdots & \tilde{D}_{mn} \end{bmatrix}$$

denote by  $\tilde{D} = (\langle l_{ij}, c_{ij}, d_{ij}, r_{ij} \rangle)_{m \times n}$  or  $\tilde{D} = (\tilde{D}_{ij})_{m \times n}$ .

From  $\tilde{D}_{ij} = \langle (l_{ij}, c_{ij}, d_{ij}, r_{ij}); t_{\tilde{D}_{ij}}, z_{\tilde{D}_{ij}} \rangle$  if  $c_{ij} = d_{ij} = p_{ij}$  ( $i = 1, 2, \dots, m; j = 1, 2, \dots, n$ ) is reduced to  $\tilde{D}_{ij} = \langle (l_{ij}, p_{ij}, r_{ij}); t_{\tilde{D}_{ij}}, z_{\tilde{D}_{ij}} \rangle$  such that  $\tilde{D}_{ij} = \langle (l_{ij}, p_{ij}, r_{ij}); t_{\tilde{D}_{ij}}, z_{\tilde{D}_{ij}} \rangle$  is call matrix games with payffs of TIFN, which is particular case of matrix games with payoffs of TrIFN.

**Definition 4.** [11] Let

$$\tilde{\nu} = \langle (\nu_1, \nu_2, \nu_3, \nu_4); t_{\tilde{\nu}}, z_{\tilde{\nu}} \rangle \quad \text{and} \quad \tilde{\omega} = \langle (\omega_1, \omega_2, \omega_3, \omega_4); t_{\tilde{\omega}}, z_{\tilde{\omega}} \rangle$$

be TrIFNs. If there are mixed strategies  $x^* \in X$  and  $y^* \in Y$  so that for any mixed strategies  $x \in X$  and  $y \in Y$  they satisfy the two conditions as follows:

- (i)  $x^{*T} \tilde{D} y \tilde{\geq} \tilde{\nu}$  and
- (ii)  $x^T \tilde{D} y^* \tilde{\leq} \tilde{\omega}$ , then  $(x^*, y^*, \tilde{\nu}, \tilde{\omega})$  is called a reasonable solution of the matrix game  $\tilde{D}$  with payoffs of TrIFN.  $x^*$  and  $y^*$  are called reasonable strategy for player I and II, respectively.  $\tilde{\nu}$  and  $\tilde{\omega}$  are called reasonable values of players I and II, respectively.

The symbol " $\tilde{\leq}$ ", " $\tilde{\geq}$ " and " $\tilde{=}$ " are an intuitionistic fuzzy version of the order relation " $\leq$ ", " $\geq$ " and " $=$ " on the real number set. The sets of all reasonable values  $\tilde{\nu}$  and  $\tilde{\omega}$  for players I and II are denoted by V and W, respectively.

**Definition 5.** [11] Assume that there exist reasonable values  $\tilde{v}^* \in V$  and  $\tilde{\omega}^* \in W$  for players I and II, respectively. If there do not exist any reasonable values  $\tilde{v} \in V (\tilde{v} \neq \tilde{v}^*)$  and  $\tilde{\omega} \in W (\tilde{\omega} \neq \tilde{\omega}^*)$  so that they satisfy the conditions as follows:

(i)  $\tilde{v} \geq \tilde{v}^*$  and,

(ii)  $\tilde{\omega} \leq \tilde{\omega}^*$ , then  $(x^*, y^*, \tilde{v}^*, \tilde{\omega}^*)$  is called a solution of the matrix game  $\tilde{D}$  with payoffs of trapezoidal intuitionistic fuzzy numbers.  $x^*$  and  $y^*$  are called the maximin strategy and minimax strategy for players I and II, respectively.  $\tilde{v}^*$  and  $\tilde{\omega}^*$  are called the gain-floor of player I and the loss-ceiling of player II, respectively.  $x^{*T} \tilde{D} y^*$  is called the value of the matrix game  $\tilde{D}$  with payoffs of TrIFN.

### 2.3 Cut sets of TrIFN

**Definition 6.** [5] A  $(\alpha, \lambda)$ -cut set of  $\tilde{A} = \langle (l, c, d, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  is a crisp subset of  $\mathbb{R}$ , which is defined as follows:

$$\tilde{A}_\alpha^\lambda = \{x | \mu_{\tilde{A}}(x) \geq \alpha, \nu_{\tilde{A}}(x) \leq \lambda\},$$

where  $0 \leq \alpha \leq t_{\tilde{A}}, z_{\tilde{A}} \leq \lambda \leq 1$  and  $0 \leq \alpha + \lambda \leq 1$ .

**Definition 7.** [5] The  $\alpha$ -cut set and  $\lambda$ -cut set of  $\tilde{A} = \langle (l, c, d, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  are a crisp subset of  $\mathbb{R}$ , which is defined as follows:

$$\tilde{A}_\alpha = \{x | \mu_{\tilde{A}}(x) \geq \alpha\}$$

and

$$\tilde{A}^\lambda = \{x | \nu_{\tilde{A}}(x) \leq \lambda\}$$

respectively.

Using the membership function of  $\tilde{A} = \langle (l, c, d, r); t_{\tilde{A}}, z_{\tilde{A}} \rangle$  and Definition 7 such that  $\tilde{A}_\alpha = \{x | \mu_{\tilde{A}}(x) \geq \alpha\}$  and  $\tilde{A}^\lambda = \{x | \nu_{\tilde{A}}(x) \leq \lambda\}$  are closed interval and calculated as follows:

$$\tilde{A}_\alpha = [L_{\tilde{A}}(\alpha), R_{\tilde{A}}(\alpha)] = \left[ \frac{(t_{\tilde{A}} - \alpha)l + \alpha c}{t_{\tilde{A}}}, \frac{(t_{\tilde{A}} - \alpha)r + \alpha d}{t_{\tilde{A}}} \right] \quad (10)$$

and

$$\tilde{A}^\lambda = [L'_{\tilde{A}}(\lambda), R'_{\tilde{A}}(\lambda)] = \left[ \frac{(1 - \lambda)c + (\lambda - z_{\tilde{A}})l}{1 - z_{\tilde{A}}}, \frac{(1 - \lambda)d + (\lambda - z_{\tilde{A}})r}{1 - z_{\tilde{A}}} \right] \quad (11)$$

respectively.

**Definition 8.** [12] Assume that the  $\alpha$ -cut set and  $\lambda$ -cut set of any TrIFNs  $\tilde{A}$  and  $\tilde{B}$  are  $\tilde{A}_\alpha = [L_{\tilde{A}}(\alpha), R_{\tilde{A}}(\alpha)]$ ,  $\tilde{A}^\lambda = [L'_{\tilde{A}}(\lambda), R'_{\tilde{A}}(\lambda)]$ ,  $\tilde{B}_\alpha = [L_{\tilde{B}}(\alpha), R_{\tilde{B}}(\alpha)]$  and  $\tilde{B}^\lambda = [L'_{\tilde{B}}(\lambda), R'_{\tilde{B}}(\lambda)]$ , respectively. Then the ranking order of the TrIFNs  $\tilde{A}$  and  $\tilde{B}$  is stipulated according to the two cases as follows:

- (i) If  $L_{\tilde{B}}(\alpha) \geq L_{\tilde{A}}(\alpha)$ ,  $R_{\tilde{B}}(\alpha) \geq R_{\tilde{A}}(\alpha)$ ,  $L'_{\tilde{B}}(\lambda) \geq L'_{\tilde{A}}(\lambda)$  and  $R'_{\tilde{B}}(\lambda) \geq R'_{\tilde{A}}(\lambda)$ , then  $\tilde{B} \geq \tilde{A}$
- (ii) If  $L_{\tilde{B}}(\alpha) \leq L_{\tilde{A}}(\alpha)$ ,  $R_{\tilde{B}}(\alpha) \leq R_{\tilde{A}}(\alpha)$ ,  $L'_{\tilde{B}}(\lambda) \leq L'_{\tilde{A}}(\lambda)$  and  $R'_{\tilde{B}}(\lambda) \leq R'_{\tilde{A}}(\lambda)$ , then  $\tilde{B} \leq \tilde{A}$ .

### 3 Mathematical programming model for the TrIFN matrix game

As the TrIFN matrix game is a zero - sum game, from Definition 3 expected payoff for player I is computed as follows:

$$\begin{aligned} E(x, y) &= x^T \tilde{D}y \\ &= \sum_{i=1}^n \sum_{j=1}^n \tilde{D}_{ij} x_i y_j \\ &= \langle (\sum_{i=1}^m \sum_{j=1}^n l_{ij} x_i y_j, \sum_{i=1}^m \sum_{j=1}^n c_{ij} x_i y_j, \sum_{i=1}^m \sum_{j=1}^n d_{ij} x_i y_j, \sum_{i=1}^m \sum_{j=1}^n r_{ij} x_i y_j); \\ &\quad \wedge \{t_{\tilde{D}_{ij}}\}, \vee \{z_{\tilde{D}_{ij}}\} \rangle \end{aligned}$$

which is a TrIFN.

And the expected payoff for player II obtained as follows:

$$\begin{aligned} E(x, y) &= x^T (-\tilde{D})y \\ &= \sum_{i=1}^n \sum_{j=1}^n (-\tilde{D}_{ij}) x_i y_j \\ &= \langle (-\sum_{i=1}^m \sum_{j=1}^n r_{ij} x_i y_j, -\sum_{i=1}^m \sum_{j=1}^n d_{ij} x_i y_j, -\sum_{i=1}^m \sum_{j=1}^n c_{ij} x_i y_j, \\ &\quad -\sum_{i=1}^m \sum_{j=1}^n l_{ij} x_i y_j); \wedge \{t_{\tilde{D}_{ij}}\}, \vee \{z_{\tilde{D}_{ij}}\} \rangle. \end{aligned}$$



Assume that player I is a maximizing player and player II is a minimizing player. Player I should choose a mixed strategy  $x \in X$  that maximizes the minimum expected gain of player II, i.e.,

$$\tilde{\nu} = \max_{x \in X} \min_{y \in Y} \{E(x, y)\}, \quad (12)$$

which is called player I's gain-floor.

Similarly, player II should choose a mixed strategy  $y \in Y$  that minimizes the maximum expected loss of player I, i.e.,

$$\tilde{\omega} = \min_{y \in Y} \max_{x \in X} \{E(x, y)\}, \quad (13)$$

which is called player II's loss-ceiling.

Hence, player I's gain-floor and player II's loss ceiling denoted by

$$\tilde{\nu} = \langle (\nu_1, \nu_2, \nu_3, \nu_4); t_{\tilde{\nu}}, z_{\tilde{\nu}} \rangle$$

and

$$\tilde{\omega} = \langle (\omega_1, \omega_2, \omega_3, \omega_4); t_{\tilde{\omega}}, z_{\tilde{\omega}} \rangle.$$

From Definitions 4, 5 and Eqs.(12) and (13) the maximin strategy  $x^*$  and gain-floor  $\tilde{\nu}^*$  of player I and the minimax strategy  $y^*$  and loss-ceiling  $\tilde{\omega}^*$  of player II can be generate by solving an intuitionistic fuzzy mathematical programming model constructed as follows:

$$\begin{aligned} & \max\{\tilde{\nu}\} \\ & s.t. \begin{cases} \sum_{i=1}^m \tilde{D}_{ij} x_i y_j \geq \tilde{\nu} & (j = 1, 2, \dots, n)(y \in Y) \\ \sum_{i=1}^m x_i = 1 \\ x_i \geq 0 & (i = 1, 2, \dots, m) \end{cases} \end{aligned} \quad (14)$$

and

$$\begin{aligned} & \min\{\tilde{\omega}\} \\ & s.t. \begin{cases} \sum_{j=1}^n \tilde{D}_{ij} x_i y_j \leq \tilde{\omega} & (i = 1, 2, \dots, m)(x \in X) \\ \sum_{j=1}^n y_j = 1 \\ y_j \geq 0 & (j = 1, 2, \dots, n), \end{cases} \end{aligned} \quad (15)$$

respectively, where  $\tilde{\nu}$  and  $\tilde{\omega}$  are TrIFNs.

From Eqs.(14), (15) and theorem [6] can be converted into an intuitionistic fuzzy mathematical programming models as follows:

$$\begin{aligned} & \max\{\tilde{\nu}\} \\ & s.t. \begin{cases} \sum_{i=1}^m \tilde{D}_{ij}x_i \geq \tilde{\nu} & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m x_i = 1 \\ x_i \geq 0 & (i = 1, 2, \dots, m) \end{cases} \end{aligned} \quad (16)$$

and

$$\begin{aligned} & \min\{\tilde{\omega}\} \\ & s.t. \begin{cases} \sum_{j=1}^n \tilde{D}_{ij}y_j \leq \tilde{\omega} & (i = 1, 2, \dots, m) \\ \sum_{j=1}^n y_j = 1 \\ y_j \geq 0 & (j = 1, 2, \dots, n) \end{cases} \end{aligned} \quad (17)$$

respectively. From Definitions 6 and 8 can be transformed into the interval-valued bi-objective mathematical programming models as follows:

$$\begin{aligned} & \max\{\tilde{\nu}_\alpha, \tilde{\nu}^\lambda\} \\ & s.t. \begin{cases} \sum_{i=1}^m (\tilde{D}_{ij})_\alpha x_i \geq \tilde{\nu}_\alpha & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m (\tilde{D}_{ij})^\lambda x_i \geq \tilde{\nu}^\lambda & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m x_i = 1 \\ x_i \geq 0 & (i = 1, 2, \dots, m) \end{cases} \end{aligned} \quad (18)$$

the  $\alpha$ -cut set and  $\lambda$ -cut set of the TrIFNs  $\tilde{\nu}$  and  $\tilde{D}_{ij} = (i = 1, 2, \dots, m; j = 1, 2, \dots, n)$  are denoted by

$$\tilde{\nu}_\alpha = [\nu_\alpha^L, \nu_\alpha^R], \quad \tilde{\nu}^\lambda = [\nu_L^\lambda, \nu_R^\lambda], \quad (\tilde{D}_{ij})_\alpha = [L_{\tilde{D}_{ij}}(\alpha), R_{\tilde{D}_{ij}}(\alpha)]$$

and

$$(\tilde{D}_{ij})^\lambda = [L'_{\tilde{D}_{ij}}(\lambda), R'_{\tilde{D}_{ij}}(\lambda)],$$

respectively. From Eq.(18) can be written as the following interval-valued bi-object mathematical programming model:

$$\begin{aligned} & \max\{[\nu_\alpha^L, \nu_\alpha^R], [\nu_L^\lambda, \nu_R^\lambda]\} \\ & s.t. \left\{ \begin{array}{l} \sum_{i=1}^m L_{\tilde{D}_{ij}}(\alpha)x_i \geq \nu_\alpha^L \quad (j = 1, 2, \dots, n) \\ \sum_{i=1}^m R_{\tilde{D}_{ij}}(\alpha)x_i \geq \nu_\alpha^R \quad (j = 1, 2, \dots, n) \\ \sum_{i=1}^m L'_{\tilde{D}_{ij}}(\lambda)x_i \geq \nu_L^\lambda \quad (j = 1, 2, \dots, n) \\ \sum_{i=1}^m R'_{\tilde{D}_{ij}}(\lambda)x_i \geq \nu_R^\lambda \quad (j = 1, 2, \dots, n) \\ \sum_{i=1}^m x_i = 1 \\ x_i \geq 0 \quad (i = 1, 2, \dots, m) \end{array} \right. \quad (19) \end{aligned}$$

where  $\nu_\alpha^L, \nu_\alpha^R, \nu_L^\lambda, \nu_R^\lambda$  and  $x_i (i = 1, 2, \dots, m)$  are decision variables.

In Eq.(19) two interval-valued objective functions, we will using the linear weighted averaging method of multiobjective decision making, their weights are the same as 1/2 hence, Eq.(19) can be aggregated into the interval-valued mathematical programming model as follows:

$$\begin{aligned}
 & \max \left\{ \left[ \frac{\nu_\alpha^L + \nu_L^\lambda}{2}, \frac{\nu_\alpha^R + \nu_R^\lambda}{2} \right] \right\} \\
 & \text{s.t.} \left\{ \begin{array}{l}
 \sum_{i=1}^m L_{\tilde{D}_{ij}}(\alpha)x_i \geq \nu_\alpha^L \quad (j = 1, 2, \dots, n) \\
 \sum_{i=1}^m R_{\tilde{D}_{ij}}(\alpha)x_i \geq \nu_\alpha^R \quad (j = 1, 2, \dots, n) \\
 \sum_{i=1}^m L'_{\tilde{D}_{ij}}(\lambda)x_i \geq \nu_L^\lambda \quad (j = 1, 2, \dots, n) \\
 \sum_{i=1}^m R'_{\tilde{D}_{ij}}(\lambda)x_i \geq \nu_R^\lambda \quad (j = 1, 2, \dots, n) \\
 \sum_{i=1}^m x_i = 1 \\
 x_i \geq 0 \quad (i = 1, 2, \dots, m).
 \end{array} \right. \quad (20)
 \end{aligned}$$

From Eq.(20) and Ishibuchi and Ianaka [2] the maximization problem with the interval- valued objective function can be written as follows:

$$\begin{aligned}
 & \max \left\{ \frac{\nu_\alpha^L + \nu_L^\lambda}{2}, \frac{\nu_\alpha^L + \nu_L^\lambda + \nu_\alpha^R + \nu_R^\lambda}{4} \right\} \\
 & \text{s.t.} \left\{ \begin{array}{l}
 \sum_{i=1}^m L_{\tilde{D}_{ij}}(\alpha)x_i \geq \nu_\alpha^L \quad (j = 1, 2, \dots, n) \\
 \sum_{i=1}^m R_{\tilde{D}_{ij}}(\alpha)x_i \geq \nu_\alpha^R \quad (j = 1, 2, \dots, n) \\
 \sum_{i=1}^m L'_{\tilde{D}_{ij}}(\lambda)x_i \geq \nu_L^\lambda \quad (j = 1, 2, \dots, n) \\
 \sum_{i=1}^m R'_{\tilde{D}_{ij}}(\lambda)x_i \geq \nu_R^\lambda \quad (j = 1, 2, \dots, n) \\
 \sum_{i=1}^m x_i = 1 \\
 x_i \geq 0 \quad (i = 1, 2, \dots, m).
 \end{array} \right. \quad (21)
 \end{aligned}$$

From Eq.(21) and using the linear weighted averaging method of multi-objective decision making [13] - [14] can be written as follows:

$$\max \left\{ \xi \frac{\nu_\alpha^L + \nu_L^\lambda}{2} + (1 - \xi) \frac{\nu_\alpha^L + \nu_L^\lambda + \nu_\alpha^R + \nu_R^\lambda}{4} \right\}$$

$$s.t. \begin{cases} \sum_{i=1}^m L_{\tilde{D}_{ij}}(\alpha) x_i \geq \nu_\alpha^L & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m R_{\tilde{D}_{ij}}(\alpha) x_i \geq \nu_\alpha^R & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m L'_{\tilde{D}_{ij}}(\lambda) x_i \geq \nu_L^\lambda & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m R'_{\tilde{D}_{ij}}(\lambda) x_i \geq \nu_R^\lambda & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m x_i = 1 \\ x_i \geq 0 & (i = 1, 2, \dots, m) \end{cases} \quad (22)$$

where  $\xi \in [0, 1]$ .

From Eqs.(10) and (11) we can obtain the  $\alpha$ -cut set and  $\lambda$ -cut set of the TrIFNs  $\tilde{D}_{ij} = \langle (l_{ij}, c_{ij}, d_{ij}, r_{ij}); t_{\tilde{D}_{ij}}, z_{\tilde{D}_{ij}} \rangle (i = 1, 2, \dots, m; j = 1, 2, \dots, n)$  as follows:

$$(\tilde{D}_{ij})_\alpha = [L_{\tilde{D}_{ij}}(\alpha), R_{\tilde{D}_{ij}}(\alpha)] = \left[ \frac{(t_{\tilde{D}_{ij}} - \alpha)l_{ij} + \alpha c_{ij}}{t_{\tilde{D}_{ij}}}, \frac{(t_{\tilde{D}_{ij}} - \alpha)r_{ij} + \alpha d_{ij}}{t_{\tilde{D}_{ij}}} \right] \quad (23)$$

and

$$\begin{aligned} (\tilde{D}_{ij})^\lambda &= [L'_{\tilde{D}_{ij}}(\lambda), R'_{\tilde{D}_{ij}}(\lambda)] \\ &= \left[ \frac{(1 - \lambda)a_{ij} + (\lambda - z_{\tilde{D}_{ij}})l_{ij}}{1 - z_{\tilde{D}_{ij}}}, \frac{(1 - \lambda)d_{ij} + (\lambda - u_{\tilde{D}_{ij}})r_{ij}}{1 - z_{\tilde{D}_{ij}}} \right] \end{aligned} \quad (24)$$

respectively.

From Eq.(22) can rewritten as the following linear programming model:

$$\begin{aligned}
 & \max \left\{ \xi \frac{\nu_\alpha^L + \nu_L^\lambda}{2} + (1 - \xi) \frac{\nu_\alpha^L + \nu_L^\lambda + \nu_\alpha^R + \nu_R^\lambda}{4} \right\} \\
 & s.t. \begin{cases} \sum_{i=1}^m \frac{(t_{\tilde{D}_{ij}} - \alpha)l_{ij} + \alpha c_{ij}}{t_{\tilde{D}_{ij}}} x_i \geq \nu_\alpha^L & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m \frac{(t_{\tilde{D}_{ij}} - \alpha)r_{ij} + \alpha d_{ij}}{t_{\tilde{D}_{ij}}} x_i \geq \nu_\alpha^R & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m \frac{(1 - \lambda)c_{ij} + (\lambda - z_{\tilde{D}_{ij}})l_{ij}}{1 - z_{\tilde{D}_{ij}}} x_i \geq \nu_L^\lambda & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m \frac{(1 - \lambda)d_{ij} + (\lambda - z_{\tilde{D}_{ij}})r_{ij}}{1 - z_{\tilde{D}_{ij}}} x_i \geq \nu_R^\lambda & (j = 1, 2, \dots, n) \\ \sum_{i=1}^m x_i = 1 \\ x_i \geq 0 & (i = 1, 2, \dots, m) \end{cases} \quad (25)
 \end{aligned}$$

where  $\xi \in [0, 1]$ ,  $\alpha \in [0, t_{\tilde{D}}]$ ,  $\lambda \in [z_{\tilde{D}}, 1]$  and  $\alpha + \lambda \in [0, 1]$ .

Using the simplex method of linear programming, we can obtain the optimal solution of Eq.(25), denote by  $(x^*(\alpha, \lambda), \nu_\alpha^{L*}, \nu_\alpha^{R*}, \nu_L^{\lambda*}, \nu_R^{\lambda*})$ . where  $x^*(\alpha, \lambda)$  is the maximin strategy of player I at the  $\langle \alpha, \lambda \rangle$ -confidence level.  $\nu_\alpha^{L*}$  and  $\nu_\alpha^{R*}$  are the lower and upper bounds of the gain-floor  $\tilde{\nu}^*$  of player I at the  $\alpha$ -confidence level, that is  $\alpha$ -cut set  $\tilde{\nu}_\alpha^*$  of  $\tilde{\nu}^*$ . Similarly,  $\nu_L^{\lambda*}$  and  $\nu_R^{\lambda*}$  are the lower and upper bounds of the gain-floor  $\tilde{\nu}^*$  of player I at the  $\lambda$ -confidence level, that is that is  $\lambda$ -cut set  $\tilde{\nu}_\lambda^*$  of  $\tilde{\nu}^*$ .

In the same way, from Definition 8 and Eq.(17) can be written as the following interval-valued bi-objective mathematical programming model:

$$\begin{aligned}
 & \min \{ \tilde{\omega}_\alpha, \tilde{\omega}^\lambda \} \\
 & s.t. \begin{cases} \sum_{j=1}^n (\tilde{D}_{ij})_\alpha y_j \leq \tilde{\omega}_\alpha & (i = 1, 2, \dots, m) \\ \sum_{j=1}^n (\tilde{D}_{ij})^\lambda y_j \leq \tilde{\omega}^\lambda & (i = 1, 2, \dots, m) \\ \sum_{j=1}^n y_j = 1 \\ y_j \geq 0 & (j = 1, 2, \dots, n). \end{cases} \quad (26)
 \end{aligned}$$

The  $\alpha$ -cut set and  $\lambda$ -cut set of the TrIFN  $\tilde{\omega}$  are denote by

$$\tilde{\omega}_\alpha = [\omega_\alpha^L, \omega_\alpha^R] \quad \text{and} \quad \tilde{\omega}^\lambda = [\omega_L^\lambda, \omega_R^\lambda],$$

respectively.

From Eq.[26] can be written as the folowing interval-valued bi-objective mathematical programming model:

$$\min\{[\omega_\alpha^L, \omega_\alpha^R], [\omega_L^\lambda, \omega_R^\lambda]\}$$

$$s.t. \begin{cases} \sum_{i=1}^n L_{\tilde{D}_{ij}}(\alpha)y_j \leq \omega_R^L & (i = 1, 2, \dots, m) \\ \sum_{i=1}^n R_{\tilde{D}_{ij}}(\alpha)y_j \leq \omega_\alpha^R & (i = 1, 2, \dots, m) \\ \sum_{i=1}^n L'_{\tilde{D}_{ij}}(\lambda)y_j \leq \omega_L^\lambda & (i = 1, 2, \dots, m) \\ \sum_{i=1}^n R'_{\tilde{D}_{ij}}(\lambda)y_j \leq \omega_R^\lambda & (i = 1, 2, \dots, m) \\ \sum_{i=1}^n y_j = 1 \\ y_j \geq 0 & (j = 1, 2, \dots, n) \end{cases} \quad (27)$$

where  $\omega_\alpha^L, \omega_\alpha^R, \omega_L^\lambda, \omega_R^\lambda$  and  $y_j$  ( $i = 1, 2, \dots, n$ ) are decision variables.

From Eq.(27) and use the linear weighted veraging method of multiobjective decisionmaking [13] - [14] can be aggregated into the interval-valued mathematical programming model as follows:

$$\begin{aligned}
 & \min \left\{ \left[ \frac{\omega_\alpha^L + \omega_L^\lambda}{2}, \frac{\omega_\alpha^R + \omega_R^\lambda}{2} \right] \right\} \\
 & s.t. \left\{ \begin{array}{l}
 \sum_{i=1}^n L_{\tilde{D}_{ij}}(\alpha) y_j \leq \omega_R^L \quad (i = 1, 2, \dots, m) \\
 \sum_{i=1}^n R_{\tilde{D}_{ij}}(\alpha) y_j \leq \omega_\alpha^R \quad (i = 1, 2, \dots, m) \\
 \sum_{i=1}^n L'_{\tilde{D}_{ij}}(\lambda) y_j \leq \omega_L^\lambda \quad (i = 1, 2, \dots, m) \\
 \sum_{i=1}^n R'_{\tilde{D}_{ij}}(\lambda) y_j \leq \omega_R^\lambda \quad (i = 1, 2, \dots, m) \\
 \sum_{i=1}^n y_j = 1 \\
 y_j \geq 0 \quad (i = 1, 2, \dots, n).
 \end{array} \right. \quad (28)
 \end{aligned}$$

From Eq.(28) and the maximization problem with the interval-valued objective function [2] can be written as follows:

$$\begin{aligned}
 & \min \left\{ \frac{\omega_\alpha^R + \omega_R^\lambda}{2} + \frac{\omega_\alpha^L + \omega_L^\lambda + \omega_\alpha^R + \omega_R^\lambda}{4} \right\} \\
 & s.t. \left\{ \begin{array}{l}
 \sum_{i=1}^n L_{\tilde{D}_{ij}}(\alpha) y_j \leq \omega_\alpha^L \quad (i = 1, 2, \dots, m) \\
 \sum_{i=1}^n R_{\tilde{D}_{ij}}(\alpha) y_j \leq \omega_\alpha^R \quad (i = 1, 2, \dots, m) \\
 \sum_{i=1}^n L'_{\tilde{D}_{ij}}(\lambda) y_j \leq \omega_L^\lambda \quad (i = 1, 2, \dots, m) \\
 \sum_{i=1}^n R'_{\tilde{D}_{ij}}(\lambda) y_j \leq \omega_R^\lambda \quad (i = 1, 2, \dots, m) \\
 \sum_{i=1}^n y_j = 1 \\
 y_j \geq 0 \quad (j = 1, 2, \dots, n).
 \end{array} \right. \quad (29)
 \end{aligned}$$

Similary, using the linear weighted averaging method of multiobjective decision making, from Eqs.(23), (24) and (29) can be further aggregated and convert into the linear programming model as follows:



$$\begin{aligned}
& \min \left\{ \xi \frac{\omega_\alpha^R + \omega_R^\lambda}{2} + (1 - \xi) \frac{\omega_\alpha^L + \omega_L^\lambda + \omega_\alpha^R + \omega_R^\lambda}{4} \right\} \\
& \text{s.t.} \left\{ \begin{array}{l}
\sum_{j=1}^n \frac{(t_{\tilde{D}_{ij}} - \alpha)l_{ij} + \alpha c_{ij}}{t_{\tilde{D}_{ij}}} y_j \leq \omega_\alpha^L \quad (i = 1, 2, \dots, m) \\
\sum_{j=1}^n \frac{(t_{\tilde{D}_{ij}} - \alpha)r_{ij} + \alpha d_{ij}}{t_{\tilde{D}_{ij}}} y_j \leq \omega_\alpha^R \quad (i = 1, 2, \dots, m) \\
\sum_{j=1}^n \frac{(1 - \lambda)c_{ij} + (\lambda - z_{\tilde{D}_{ij}})l_{ij}}{1 - z_{\tilde{D}_{ij}}} y_i \leq \omega_L^\lambda \quad (i = 1, 2, \dots, m) \\
\sum_{j=1}^n \frac{(1 - \lambda)d_{ij} + (\lambda - z_{\tilde{D}_{ij}})r_{ij}}{1 - z_{\tilde{D}_{ij}}} y_i \leq \omega_R^\lambda \quad (i = 1, 2, \dots, m) \\
\sum_{j=1}^n y_j = 1 \\
y_j \geq 0 \quad (j = 1, 2, \dots, n).
\end{array} \right. \quad (30)
\end{aligned}$$

For any adequately given values of the parameters  $\xi$ ,  $\alpha$  and  $\lambda$ , using the simplex method of linear programming, we can obtain the optimal solution of Eq.(30), denote by  $(y^*(\alpha, \lambda), \omega_\alpha^{L*}, \omega_\alpha^{R*}, \omega_L^{\lambda*}, \omega_R^{\lambda*})$  where  $y^*(\alpha, \lambda)$  is the min-max strategy of player II at the  $\langle \alpha, \lambda \rangle$ -confidence level.  $\omega_\alpha^{L*}$  and  $\omega_\alpha^{R*}$  are the lower and upper bounds of the gain-floor  $\tilde{\omega}^*$  of player II at the  $\alpha$ -confidence level, that is  $\alpha$ - cut set  $\tilde{\omega}_\alpha^*$  of  $\tilde{\omega}^*$ . similarly,  $\omega_L^{\lambda*}$  and  $\omega_R^{\lambda*}$  are the lower and upper bounds of the gain-floor  $\tilde{\omega}^*$  of player II at the  $\lambda$ -confidence level, that is that is  $\lambda$ - cut set  $\tilde{\omega}_\lambda^*$  of  $\tilde{\omega}^*$ .

#### 4 An application to voting share problem

In this examples 1 and 2, using Eqs.(25) and (30) to solve the problem.

##### Example 1.

Assume that there is an election where two major political parties M and W participate and total number of voters in that regions is stable. It means that the increase in percentage of votes for one political party results in the same for the other political party. Suppose M has two strategies as

$\rho_1$ : the campaign by big rallies and superstar.

$\rho_2$ : co-operating with other small political parties to reduce secured votes of the opposition.

Simultaneously W takes two strategies:

$\tau_1$ : Making lot of promises to the people.

$\tau_2$ : campaigning by use of mixed media such as publication and television.

Let us consider matrix game D with payoffs of TrIFN, where the payoff matrix of the political parties M is given as follows:

$$\tilde{D} = \begin{bmatrix} \langle(155, 165, 175, 180); 0.7, 0.2\rangle & \langle(130, 146, 150, 165); 0.6, 0.2\rangle \\ \langle(75, 85, 95, 100); 0.6, 0.3\rangle & \langle(160, 170, 184, 190); 0.8, 0.1\rangle \end{bmatrix}$$

where the element  $\langle(155, 165, 175, 180); 0.7, 0.2\rangle$  in the matrix  $\tilde{D}$  is TrIFN represent that when M choose the strategy  $\rho_1$  and W choose the strategy  $\tau_1$  then votes of the political parties M is between 155 and 180 the maximum confidence level and minimum non-confidence level of the head election exponent of  $\tilde{D}$  are 0.7 and 0.2, respectively. In this case, the hestiance degree is 0.1 other elements in  $\tilde{D}$  may be identically explained.

From eqs.(25) and (30) the parameterized linear programming is obtained as follows:

$$\begin{aligned} & \max \left\{ \xi \frac{\nu_\alpha^L + \nu_L^\lambda}{2} + (1 - \xi) \frac{\nu_\alpha^L + \nu_L^\lambda + \nu_\alpha^R + \nu_R^\lambda}{4} \right\} \\ & s.t. \left\{ \begin{array}{l} \frac{(0.7 - \alpha)155 + 165\alpha}{0.7}x_1 + \frac{(0.6 - \alpha)75 + 85\alpha}{0.6}x_2 \geq \nu_\alpha^L \\ \frac{(0.6 - \alpha)130 + 146\alpha}{0.6}x_1 + \frac{(0.8 - \alpha)160 + 170\alpha}{0.8}x_2 \geq \nu_\alpha^L \\ \frac{(0.7 - \alpha)180 + 175\alpha}{0.6}x_1 + \frac{(0.6 - \alpha)100 + 95\alpha}{0.6}x_2 \geq \nu_\alpha^R \\ \frac{(0.6 - \alpha)165 + 150\alpha}{0.7}x_1 + \frac{(0.8 - \alpha)190 + 184\alpha}{0.8}x_2 \geq \nu_\alpha^R \\ \frac{(1 - \lambda)165 + (\lambda - 0.2)155}{0.6}x_1 + \frac{(1 - \lambda)85 + (\lambda - 0.3)75}{0.8}x_2 \geq \nu_L^\lambda \\ \frac{(1 - \lambda)146 + (\lambda - 0.2)130}{0.8}x_1 + \frac{(1 - \lambda)170 + (\lambda - 0.1)160}{0.7}x_2 \geq \nu_L^\lambda \\ \frac{(1 - \lambda)175 + (\lambda - 0.2)180}{0.8}x_1 + \frac{(1 - \lambda)95 + (\lambda - 0.3)100}{0.9}x_2 \geq \nu_R^\lambda \\ \frac{(1 - \lambda)150 + (\lambda - 0.2)165}{0.8}x_1 + \frac{(1 - \lambda)184 + (\lambda - 0.1)190}{0.9}x_2 \geq \nu_R^\lambda \\ x_1 + x_2 = 1 \\ x_1 \geq 0, x_2 \geq 0 \end{array} \right. \end{aligned} \quad (31)$$

and

$$\min \left\{ \xi \frac{\omega_\alpha^R + \omega_R^\lambda}{2} + (1 - \xi) \frac{\omega_\alpha^L + \omega_L^\lambda + \omega_\alpha^R + \omega_R^\lambda}{4} \right\}$$

$$s.t. \left\{ \begin{array}{l} \frac{(0.7 - \alpha)155 + 165\alpha}{(0.6 - \alpha)^{0.7} 75 + 85\alpha} y_1 + \frac{(0.6 - \alpha)130 + 146\alpha}{(0.8 - \alpha)^{0.6} 160 + 170\alpha} y_2 \leq \omega_\alpha^L \\ \frac{(0.7 - \alpha)^{0.6} 180 + 175\alpha}{(0.6 - \alpha)^{0.8} 165 + 150\alpha} y_1 + \frac{(0.6 - \alpha)^{0.6} 190 + 184\alpha}{(1 - \lambda)165 + (\lambda - 0.2)155} y_2 \leq \omega_\alpha^R \\ \frac{(0.6 - \alpha)^{0.7} 100 + 95\alpha}{(1 - \lambda)85 + (\lambda - 0.3)75} y_1 + \frac{(0.8 - \alpha)^{0.6} 190 + 184\alpha}{(1 - \lambda)170 + (\lambda - 0.1)160} y_2 \leq \omega_\alpha^R \\ \frac{(1 - \lambda)165 + (\lambda - 0.2)155}{(1 - \lambda)175 + (\lambda - 0.2)180} y_1 + \frac{(1 - \lambda)146 + (\lambda - 0.2)130}{(1 - \lambda)150 + (\lambda - 0.2)165} y_2 \leq \omega_L^\lambda \\ \frac{(1 - \lambda)85 + (\lambda - 0.3)75}{(1 - \lambda)95 + (\lambda - 0.3)100} y_1 + \frac{(1 - \lambda)170 + (\lambda - 0.1)160}{(1 - \lambda)184 + (\lambda - 0.1)190} y_2 \leq \omega_L^\lambda \\ \frac{(1 - \lambda)175 + (\lambda - 0.2)180}{(1 - \lambda)95 + (\lambda - 0.3)100} y_1 + \frac{(1 - \lambda)150 + (\lambda - 0.2)165}{(1 - \lambda)184 + (\lambda - 0.1)190} y_2 \leq \omega_R^\lambda \\ \frac{(1 - \lambda)95 + (\lambda - 0.3)100}{0.7} y_1 + \frac{(1 - \lambda)184 + (\lambda - 0.1)190}{0.9} y_2 \leq \omega_R^\lambda \\ y_1 + y_2 = 1 \\ y_1 \geq 0, y_2 \geq 0 \end{array} \right. \quad (32)$$

respectively.

For the given  $\xi = 0.8$ ,  $\alpha \in [0, 0.6]$  and  $\lambda \in [0.3, 1]$ . The greatest possible value of  $\alpha$  and the smallest possible value of  $\lambda$  are computed as follows:

$$\min\{t_{\tilde{D}_{ij}} | i = 1, 2; J = 1, 2\} = \min\{0.7, 0.6, 0.6, 0.8\} = 0.6$$

and

$$\max\{z_{\tilde{D}_{ij}} | i = 1, 2; J = 1, 2\} = \max\{0.2, 0.2, 0.3, 0.1\} = 0.3$$

respectively.

Solving Eqs.(31) and (32) we computed by simplex method for linear programming. The result as follows:

Player M's gain-floor  $\tilde{v}^*$  stay in the ranges  $\tilde{v}_{(\alpha, \lambda)}^* = [136.82, 161.82]$ . When  $\langle \alpha, \lambda \rangle = \langle 0.6, 0.3 \rangle$ ,  $\tilde{v}_{(\alpha, \lambda)}^* = [148.18, 157.75]$  is the most possible value of the gain-floor  $\tilde{v}^*$  of player M, where  $x^{*T}(\alpha, \lambda) = (0.824, 0.176)$ . And Player W's

loss-ceiling  $\tilde{\omega}^*$  stay in the ranges  $\tilde{\omega}_{\langle\alpha,\lambda\rangle}^* = [139.76, 168.57]$ . When  $\langle\alpha, \lambda\rangle = \langle 0.6, 0.3\rangle$ ,  $\tilde{\omega}_{\langle\alpha,\lambda\rangle}^* = [150.05, 159.13]$  is the most possible value of the loss-ceiling  $\tilde{\omega}^*$  of player W, where  $y^{*T}(\alpha, \lambda) = (0.306, 0.694)$ . Thus, the approximate values of player M's gain-floor  $\tilde{\nu}^*$  and player W's loss-ceiling  $\tilde{\omega}^*$  which are TrIFNs are  $\tilde{\nu}^* = \langle(136.82, 148.18, 157.75, 161.82); 0.6, 0.3\rangle$  and  $\tilde{\omega}^* = \langle(139.76, 150.05, 159.13, 168.57); 0.6, 0.3\rangle$ , respectively.

**Example 2.**

From example 1 we will reduced matrix game  $\tilde{D}$  with payoffs of TrIFN to matrix game  $\tilde{D}$  of TIFN as follows:

$$\tilde{D} = \begin{bmatrix} \langle(155, 170, 180); 0.7, 0.2\rangle & \langle(130, 148, 165); 0.6, 0.2\rangle \\ \langle(75, 90, 100); 0.6, 0.3\rangle & \langle(160, 177, 190); 0.8, 0.1\rangle \end{bmatrix}$$

From eqs.(25) and (30) the parameterized linear programming is obtained as follows:

$$\begin{aligned} & \max \left\{ \xi \frac{\nu_\alpha^L + \nu_L^\lambda}{2} + (1 - \xi) \frac{\nu_\alpha^L + \nu_L^\lambda + \nu_\alpha^R + \nu_R^\lambda}{4} \right\} \\ & s.t. \left\{ \begin{array}{l} \frac{(0.7 - \alpha)155 + 170\alpha}{(0.6 - \alpha)130 + 148\alpha} x_1 + \frac{(0.6 - \alpha)75 + 90\alpha}{(0.8 - \alpha)160 + 177\alpha} x_2 \geq \nu_\alpha^L \\ \frac{0.7}{(0.6 - \alpha)130 + 148\alpha} x_1 + \frac{0.6}{(0.8 - \alpha)160 + 177\alpha} x_2 \geq \nu_\alpha^L \\ \frac{0.6}{(0.7 - \alpha)180 + 170\alpha} x_1 + \frac{0.8}{(0.6 - \alpha)100 + 90\alpha} x_2 \geq \nu_\alpha^R \\ \frac{0.7}{(0.6 - \alpha)165 + 148\alpha} x_1 + \frac{0.6}{(0.8 - \alpha)190 + 177\alpha} x_2 \geq \nu_\alpha^R \\ \frac{0.6}{(1 - \lambda)170 + (\lambda - 0.2)155} x_1 + \frac{0.8}{(1 - \lambda)90 + (\lambda - 0.3)75} x_2 \geq \nu_L^\lambda \\ \frac{0.8}{(1 - \lambda)148 + (\lambda - 0.2)130} x_1 + \frac{0.7}{(1 - \lambda)177 + (\lambda - 0.1)160} x_2 \geq \nu_L^\lambda \\ \frac{0.8}{(1 - \lambda)170 + (\lambda - 0.2)180} x_1 + \frac{0.9}{(1 - \lambda)90 + (\lambda - 0.3)100} x_2 \geq \nu_R^\lambda \\ \frac{0.8}{(1 - \lambda)148 + (\lambda - 0.2)165} x_1 + \frac{0.7}{(1 - \lambda)177 + (\lambda - 0.1)190} x_2 \geq \nu_R^\lambda \\ x_1 + x_2 = 1 \\ x_1 \geq 0, x_2 \geq 0 \end{array} \right. \end{aligned} \tag{33}$$

and

$$\min \left\{ \xi \frac{\omega_\alpha^R + \omega_R^\lambda}{2} + (1 - \xi) \frac{\omega_\alpha^L + \omega_L^\lambda + \omega_\alpha^R + \omega_R^\lambda}{4} \right\}$$

$$s.t. \left\{ \begin{array}{l} \frac{(0.7 - \alpha)155 + 170\alpha}{0.7} y_1 + \frac{(0.6 - \alpha)130 + 148\alpha}{0.6} y_2 \leq \omega_\alpha^L \\ \frac{(0.6 - \alpha)75 + 90\alpha}{0.6} y_1 + \frac{(0.8 - \alpha)160 + 177\alpha}{0.8} y_2 \leq \omega_\alpha^L \\ \frac{(0.7 - \alpha)180 + 170\alpha}{0.6} y_1 + \frac{(0.6 - \alpha)165 + 148\alpha}{0.8} y_2 \leq \omega_\alpha^R \\ \frac{(0.6 - \alpha)100 + 90\alpha}{0.7} y_1 + \frac{(0.8 - \alpha)190 + 177\alpha}{0.6} y_2 \leq \omega_\alpha^R \\ \frac{(1 - \lambda)170 + (\lambda - 0.2)155}{0.8} y_1 + \frac{(1 - \lambda)148 + (\lambda - 0.2)130}{0.8} y_2 \leq \omega_L^\lambda \\ \frac{(1 - \lambda)90 + (\lambda - 0.3)75}{0.8} y_1 + \frac{(1 - \lambda)177 + (\lambda - 0.1)160}{0.8} y_2 \leq \omega_L^\lambda \\ \frac{(1 - \lambda)170 + (\lambda - 0.2)180}{0.8} y_1 + \frac{(1 - \lambda)148 + (\lambda - 0.2)165}{0.9} y_2 \leq \omega_R^\lambda \\ \frac{(1 - \lambda)90 + (\lambda - 0.3)100}{0.7} y_1 + \frac{(1 - \lambda)177 + (\lambda - 0.1)190}{0.9} y_2 \leq \omega_R^\lambda \\ y_1 + y_2 = 1 \\ y_1 \geq 0, y_2 \geq 0 \end{array} \right. \quad (34)$$

respectively.

Solving Eqs.(33) and (34) we computed by simplex method for linear programming. The result as follows: Player M's gain-floor  $\tilde{v}^*$  stay in the ranges  $\tilde{v}_{(\alpha, \lambda)}^* = [136.82, 161.82]$ . When  $\langle \alpha, \lambda \rangle = \langle 0.6, 0.3 \rangle$ ,  $\tilde{v}_{(\alpha, \lambda)}^* = 152.22$  is the most possible value of the gain-floor  $\tilde{v}^*$  of player M, where  $x^{*T}(\alpha, \lambda) = (0.806, 0.194)$ . And Player W's loss-ceiling  $\tilde{\omega}^*$  stay in the ranges  $\tilde{\omega}_{(\alpha, \lambda)}^* = [139.76, 168.57]$ . When  $\langle \alpha, \lambda \rangle = \langle 0.6, 0.3 \rangle$ ,  $\tilde{\omega}_{(\alpha, \lambda)}^* = 155.13$  is the most possible value of the loss-ceiling  $\tilde{\omega}^*$  of player W, where  $y^{*T}(\alpha, \lambda) = (0.284, 0.716)$ . Thus, the approximate values of player M's gain-floor  $\tilde{v}^*$  and player W's loss-ceiling  $\tilde{\omega}^*$  which are TrIFNs are  $\tilde{v}^* = \langle (136.82, 152.22, 161.82); 0.6, 0.3 \rangle$  and  $\tilde{\omega}^* = \langle (139.76, 155.13, 168.57); 0.6, 0.3 \rangle$ , respectively.

From examples 1 and 2 we will find the estimate values of player M's gain-floor  $\tilde{v}^*$  and player W's loss-ceiling  $\tilde{\omega}^*$  for matrix game with payoffs of TIFN and matrix game with payoffs of TrIFN are a little different. The matrix game with payoffs of TIFN is the most possible single value, while the matrix game with payoffs of TrIFN is the range the most possible.

## 5 Conclusion

Game theory is about the strategy of the decision of player. Good decisions require accurate and precise data. But in some cases the information available to a fuzzy uncertainty will affect the decision and payoffs.

In this work we have examples of two-person zero-sum games which only two players who have defined the strategy of each player on two strategies. To find the best response we have used the concept of the cut sets and concept of solution of matrix games with payoffs of TrIFNs. Intuitionistic fuzzy linear programming models are established for two players, which change into bi-objective parameterized linear programming model. Two linear programming models are constructed to generate the maximin and minimax strategies for players it is seen the solving TrIFN matrix games be comes to solving a pair of intuitionistic fuzzy linear programming problems. This is one example with respect to the election. We also can apply to the issue of market share, inventory management[15], [16], finance[17], management and economics.

**Competing interests:** The authors declare that they have no competing interests.

**Authors' contributions:** All authors have equal contributions. All authors read and approved final manuscript.

**Acknowledgements:** This project was supported by the Theoretical and Computational Science (TaCS) Center (Project Grant No.TaCS2559-1) under Computational and Applied Science for Smart Innovation Research Cluster (CLASSIC), Faculty of Science, KMUTT.

## References

- [1] K.T. Atanassov. Intuitionistic fuzzy sets. *Fuzzy set Syst.*, 20(1):87–96, 1986. doi: 10.1016/S0165-0114(86)80034-3.
- [2] H. Ishibuchi and H. Tanaka. Multiobjective programming in optimization of the interval objective function. *Eur. J. Oper. Res.*, 48(2):219–225, 1990. doi: 10.1016/0377-2217(90)90375-L.

- 
- [3] S. Chanas and D. Kuchta. Multiobjective programming in optimization of interval objective functions - a generalized approach. *European Journal of Operational Research*, 94(1):594–598, 1996. doi: 10.1016/0377-2217(95)00055-0.
- [4] A.C. Cevikel and M. Ahlatçoglu. Solutions for fuzzy matrix games. *Computers and Mathematics with Applications*, 60(3):399–410, 2010. doi: /10.1016/j.camwa.2010.04.020.
- [5] J.X. Nan, D.F. Li, and M.J. Zhang. A lexicographic method for matrix games with payoffs of triangular intuitionistic fuzzy numbers. *International Journal of Computational Intelligence Systems*, 3(3):280–289, 2010. doi: 10.1080/18756891.2010.9727699.
- [6] J.X. Nan, M.J. Zhang, and D.F. Li. Intuitionistic fuzzy programming model for matrix games with payoffs or trapezoidal. *International Journal of Fuzzy System*, 16(4):444–456, 2014. [http://www.ijfs.org.tw/ePublication/2014\\_paper\\_4/ijfs16-4-r-2-20140829180853\\_v2.pdf](http://www.ijfs.org.tw/ePublication/2014_paper_4/ijfs16-4-r-2-20140829180853_v2.pdf).
- [7] L. Campos. Fuzzy linear programming models to solve fuzzy matrix games. *Fuzzy Set and Systems*, 32(3):257–289, 1989. doi: 10.1016/0165-0114(89)90260-1.
- [8] D. Dubey, S. Chandra, and A. Mehra. Fuzzy linear programming under interval uncertainty based on IFS representation. *Fuzzy set and Systems*, 188(1):68–87, 2012. doi: 10.1016/j.fss.2011.09.008.
- [9] A. Aggarwal, S. Chandra, and A. Mehra. Solving matrix games with i-fuzzy payoffs: Pareto-optimal security strategies approach. *Fuzzy information and engineering*, pages 167–192, 2014. doi: 10.1016/j.fiae.2014.08.003.
- [10] D. Dubois and H. Prade. *Fuzzy set and Systems Theory and Application*. Academic Press, New York, 1980. ISBN: 978-0122227509.
- [11] C.R. Bector and S. Chandra. *Fuzzy mathematical programming and fuzzy matrix games. Studies in fuzziness and soft computing, Vol.169*. Springer-Verlag Berlin Heidelberg, Berlin, 2005. ISBN: 978-3-540-32371-6.

- 
- [12] D.F. Li. *Decision and Game Theory Management with Intuitionistic Fuzzy Sets*. Springer-Verlag Berlin Heidelberg, New York, 2014. ISBN: 978-3-642-40712-3.
- [13] D.F. Li. *Fuzzy Multiobjective Many Person Decision Makings and Games*. National Defense Industry Press, Beijing, 2003.
- [14] V. Chankong and Y.Y. Haimes. *Multiobjective Decision Making. Theory and Methodology (North Holland series in system science and engineering)*. Elsevier Science Ltd, New York, 1983. ISBN: 978-0444007100.
- [15] Vandana and B.K. Sharma. An inventory model for non-instantaneous deteriorating items with quadratic demand rate and shortages under trade credit policy. *Journal of Applied Analysis and Computation*, 6(3):720–737, 2016. doi: 10.11948/2016047.
- [16] Vandana and B.K. Sharma. An eoq model for retailers partial permissible delay in payment linked to order quantity with shortages. *Math. Comput. Simulation*, 125(1):99–112, 2016. doi: 10.1016/j.matcom.2015.11.008.
- [17] L.N. Mishra, R.P. Agarwal, and M. Sen. Solvability and asymptotic behavior for some nonlinear quadratic integral equation involving erdélyi-kober fractional integrals on the unbounded interval. *Progress in Fraction Differentiation and Applications*, 2(3):153–168, 2016. doi: 10.18576/pfda/020301.